

FEKETE-SZEGÖ PROBLEM AND SECOND HANKEL DETERMINANT FOR A CLASS OF BI-UNIVALENT FUNCTIONS

N. MAGESH AND J. YAMINI

ABSTRACT. In this sequel to the recent work (see Azizi et al., 2015), we investigate a subclass of analytic and bi-univalent functions in the open unit disk. We obtain bounds for initial coefficients, the Fekete-Szegő inequality and the second Hankel determinant inequality for functions belonging to this subclass. We also discuss some new and known special cases, which can be deduced from our results.

1. INTRODUCTION

Let \mathcal{A} denote the class of functions of the form

$$f(z) = z + \sum_{n=2}^{\infty} a_n z^n \quad (1.1)$$

which are analytic in the open unit disc $\mathbb{U} = \{z : z \in \mathbb{C} \text{ and } |z| < 1\}$ and let \mathcal{S} denote the class of functions in \mathcal{A} that are univalent in \mathbb{U} .

For two functions f and g , analytic in \mathbb{U} , we say that the function f is subordinate to g in \mathbb{U} , and write $f \prec g$, if there exists a Schwarz function w , analytic in \mathbb{U} , with $w(0) = 0$ and $|w(z)| < 1$ such that $f(z) = g(w(z))$; $z, w \in \mathbb{U}$. In particular, if the function g is univalent in \mathbb{U} , the above subordination is equivalent to $f(0) = g(0)$ and $f(\mathbb{U}) \subset g(\mathbb{U})$.

Let φ be an analytic and univalent function with positive real part in \mathbb{U} , $\varphi(0) = 1$, $\varphi'(0) > 0$ and φ maps the unit disk \mathbb{U} onto a region starlike with respect to 1 and symmetric with respect to the real axis. The Taylor's series expansion of such function is

$$\varphi(z) = 1 + B_1 z + B_2 z^2 + B_3 z^3 + \dots, \quad (1.2)$$

where all coefficients are real and $B_1 > 0$. Throughout this paper we assume that the function φ satisfies the above conditions unless otherwise stated.

By $\mathcal{S}^*(\varphi)$ and $\mathcal{K}(\varphi)$ we denote the following classes:

$$\mathcal{S}^*(\varphi) := \left\{ f \in \mathcal{S} : \frac{zf'(z)}{f(z)} \prec \varphi(z); z \in \mathbb{U} \right\}$$

and

$$\mathcal{K}(\varphi) := \left\{ f \in \mathcal{S} : 1 + \frac{zf''(z)}{f'(z)} \prec \varphi(z); z \in \mathbb{U} \right\}.$$

* Corresponding author : N. Magesh (nmagi_2000@yahoo.co.in).

The authors would like to thank Prof. H. Orhan, Department of Mathematics, Faculty of Science, Ataturk University, 25240 Erzurum, Turkey for his guidance and support.

2010 Mathematics Subject Classification : 30C45; 30C50.

Keywords and Phrases: Bi-univalent functions, bi-convex functions, Fekete-Szegő inequalities, Hankel determinants.

The classes $\mathcal{S}^*(\varphi)$ and $\mathcal{K}(\varphi)$ are the extensions of a classical set of starlike and convex functions (e.g. see Ma and Minda [20]). For $0 \leq \beta < 1$, the classes $\mathcal{S}^*(\beta) := \mathcal{S}^*\left(\frac{1+(1-2\beta)z}{1+z}\right)$ and $\mathcal{K}(\beta) := \mathcal{K}\left(\frac{1+(1-2\beta)z}{1+z}\right)$ are starlike and convex functions of order β .

It is well known (e.g. see Duren [12]) that every function $f \in \mathcal{S}$ has an inverse map f^{-1} , defined by $f^{-1}(f(z)) = z$, $z \in \mathbb{U}$ and $f(f^{-1}(w)) = w$, ($|w| < r_0(f)$; $r_0(f) \geq \frac{1}{4}$), where

$$f^{-1}(w) = w - a_2 w^2 + (2a_2^2 - a_3)w^3 - (5a_2^3 - 5a_2 a_3 + a_4)w^4 + \dots \quad (1.3)$$

A function $f \in \mathcal{A}$ is said to be bi-univalent in \mathbb{U} if both f and f^{-1} are univalent in \mathbb{U} . We let σ denote the class of bi-univalent functions in \mathbb{U} given by (1.1). A function f is said to be bi-starlike of Ma-Minda type or bi-convex of Ma-Minda type if both f and f^{-1} are, respectively, of Ma-Minda starlike or convex type. These classes are denoted, respectively, by $\mathcal{S}_\sigma^*(\varphi)$ and $\mathcal{K}_\sigma(\varphi)$ (see [3]). For $0 \leq \beta < 1$, a function $f \in \sigma$ is in the class $\mathcal{S}_\sigma^*(\beta)$ of bi-starlike functions of order β , or $\mathcal{K}_\sigma(\beta)$ of bi-convex functions of order β if both f and its inverse map f^{-1} are, respectively, starlike or convex of order β . For a history and examples of functions which are (or which are not) in the class σ , together with various other properties of subclasses of bi-univalent functions one can refer [3, 6, 7, 14, 22, 24, 28, 29].

For integers $n \geq 1$ and $q \geq 1$, the q -th Hankel determinant, defined as

$$H_q(n) = \begin{vmatrix} a_n & a_{n+1} & \cdots & a_{n+q-1} \\ a_{n+1} & a_{n+2} & \cdots & a_{n+q-2} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n+q-1} & a_{n+q-2} & \cdots & a_{n+2q-2} \end{vmatrix} \quad (a_1 = 1).$$

The Hankel determinant plays an important role in the study of singularities (see [11]). This is also an important in the study of power series with integral coefficients [8, 11]. The Hankel determinants $H_2(1) = a_3 - a_2^2$ and $H_2(2) = a_2 a_4 - a_3^2$ are well-known as Fekete-Szegő and second Hankel determinant functionals respectively. Further Fekete and Szegő [13] introduced the generalized functional $a_3 - \delta a_2^2$, where δ is some real number. In 1969, Keogh and Merkes [18] discussed the Fekete-Szegő problem for the classes starlike and convex functions. Recently, several authors have investigated upper bounds for the Hankel determinant of functions belonging to various subclasses of univalent functions [2, 9, 19, 21] and the references therein. On the other hand, Zaprawa [29, 30] extended the study of Fekete-Szegő problem to certain subclasses of bi-univalent function class σ . Following Zaprawa [29, 30], the Fekete-Szegő problem for functions belonging to various other subclasses of bi-univalent functions were considered in [17, 23]. Very recently, the upper bounds of $H_2(2)$ for the classes $\mathcal{S}_\sigma^*(\beta)$ and $\mathcal{K}_\sigma(\beta)$ were discussed by Deniz et al. [10]. Recently, Lee et al. [19] introduced the following class:

$$\mathcal{G}^\lambda(\varphi) := \left\{ f \in \mathcal{S} : (1-\lambda)f'(z) + \lambda \left(1 + \frac{zf''(z)}{f'(z)} \right) \prec \varphi(z); z \in \mathbb{U} \right\}$$

and obtained the bound for the second Hankel determinant of functions in $\mathcal{G}^\lambda(\varphi)$. It is interesting to note that

$$\mathcal{G}^\lambda := \mathcal{G}^\lambda \left(\frac{1+z}{1-z} \right) = \left\{ f : f \in \mathcal{S} \text{ and } \Re \left((1-\lambda)f'(z) + \lambda \left(1 + \frac{zf''(z)}{f'(z)} \right) \right) > 0; z \in \mathbb{U} \right\}.$$

The class \mathcal{G}^λ introduced by Al-Amiri and Reade [1]. The univalence of the functions in the class \mathcal{G}^λ was investigated by Singh et al. [26, 27].

Motivated by the recent publications (especially [4, 10, 17, 24, 29, 30]), we define the following subclass of σ .

Definition 1.1. For $0 \leq \lambda \leq 1$ and $0 \leq \beta < 1$, a function $f \in \sigma$ given by (1.1) is said to be in the class $\mathcal{G}_\sigma^\lambda(\varphi)$ if the following conditions are satisfied:

$$(1 - \lambda)f'(z) + \lambda \left(1 + \frac{zf''(z)}{f'(z)} \right) \prec \varphi(z), \quad 0 \leq \lambda \leq 1, z \in \mathbb{U}$$

and for $g = f^{-1}$ given by (1.3)

$$(1 - \lambda)g'(w) + \lambda \left(1 + \frac{wg''(w)}{g'(w)} \right) \prec \varphi(w), \quad 0 \leq \lambda \leq 1, w \in \mathbb{U}.$$

From among the many choices of φ and λ which would provide the following known subclasses:

- (1) $\mathcal{G}_\sigma^0(\varphi) := \mathcal{H}_\sigma(\varphi)$ [3],
- (2) $\mathcal{G}_\sigma^1(\varphi) := \mathcal{K}_\sigma(\varphi)$ [3],
- (3) $\mathcal{G}_\sigma^\lambda\left(\frac{1+(1-2\beta)z}{1-z}\right) := \mathcal{G}_\sigma^\lambda(\beta)$ ($0 \leq \beta < 1$) [4].
- (4) $\mathcal{G}_\sigma^0\left(\frac{1+(1-2\beta)z}{1-z}\right) := \mathcal{H}_\sigma^\beta$ ($0 \leq \beta < 1$) [28]
- (5) $\mathcal{G}_\sigma^1\left(\frac{1+(1-2\beta)z}{1-z}\right) := \mathcal{K}_\sigma(\beta)$ ($0 \leq \beta < 1$) [5].

In this paper we shall obtain the Fekete-Szegő inequalities for $\mathcal{G}_\sigma^\lambda(\varphi)$ as well as its special classes. Further, the second Hankel determinant obtained for the class $\mathcal{G}_\sigma^\lambda(\beta)$.

2. INITIAL COEFFICIENT BOUNDS

Theorem 2.1. If f given by (1.1) is in the class $\mathcal{G}_\sigma^\lambda(\varphi)$, then

$$|a_2| \leq \frac{B_1 \sqrt{B_1}}{\sqrt{4B_1 + |(3 - \lambda)B_1^2 - 4B_2|}} \quad (2.1)$$

and

$$|a_3| \leq \begin{cases} \left(1 - \frac{4}{3(1+\lambda)B_1} \right) \frac{B_1^3}{4B_1 + |(3-\lambda)B_1^2 - 4B_2|} + \frac{B_1}{3(1+\lambda)}, & \text{if } B_1 \geq \frac{4}{3(1+\lambda)}; \\ \frac{B_1}{3(1+\lambda)}, & \text{if } B_1 < \frac{4}{3(1+\lambda)}. \end{cases} \quad (2.2)$$

Proof. Suppose that $u(z)$ and $v(z)$ are analytic in the unit disk \mathbb{U} with $u(0) = v(0) = 0$, $|u(z)| < 1$, $|v(z)| < 1$ and

$$u(z) = b_1 z + \sum_{n=2}^{\infty} b_n z^n, \quad v(z) = c_1 z + \sum_{n=2}^{\infty} c_n z^n, \quad |z| < 1. \quad (2.3)$$

It is well known that

$$|b_1| \leq 1, |b_2| \leq 1 - |b_1|^2, |c_1| \leq 1, |c_2| \leq 1 - |c_1|^2. \quad (2.4)$$

By a simple calculation, we have

$$\varphi(u(z)) = 1 + B_1 b_1 z + (B_1 b_2 + B_2 b_1^2) z^2 + \dots, \quad |z| < 1 \quad (2.5)$$

and

$$\varphi(v(w)) = 1 + B_1 c_1 w + (B_1 c_2 + B_2 c_1^2) w^2 + \dots, \quad |w| < 1. \quad (2.6)$$

Let $f \in \mathcal{G}_\sigma^\lambda(\varphi)$. Then there are analytic functions $u, v : \mathbb{U} \rightarrow \mathbb{U}$ given by (2.3) such that

$$(1 - \lambda)f'(z) + \lambda \left(1 + \frac{zf''(z)}{f'(z)} \right) = \varphi(u(z)) \quad (2.7)$$

and

$$(1 - \lambda)g'(w) + \lambda \left(1 + \frac{wg''(w)}{g'(w)} \right) = \varphi(v(w)). \quad (2.8)$$

It follows from (2.5), (2.6), (2.7) and (2.8) that

$$2a_2 = B_1b_1 \quad (2.9)$$

$$3(1 + \lambda)a_3 - 4\lambda a_2^2 = B_1b_2 + B_2b_1^2 \quad (2.10)$$

$$-2a_2 = B_1c_1 \quad (2.11)$$

$$2(\lambda + 3)a_2^2 - 3(1 + \lambda)a_3 = B_1c_2 + B_2c_1^2. \quad (2.12)$$

From (2.9) and (2.11), we get

$$b_1 = -c_1. \quad (2.13)$$

By adding (2.10) to (2.12), further, using (2.9) and (2.13), we have

$$(2(3 - \lambda)B_1^2 - 8B_2)a_2^2 = B_1^3(b_2 + c_2). \quad (2.14)$$

In view of (2.13) and (2.14), together with (2.4), we get

$$|(2(3 - \lambda)B_1^2 - 8B_2)a_2^2| \leq 2B_1^3(1 - |b_1|^2). \quad (2.15)$$

Substituting (2.9) in (2.15) we obtain

$$|a_2| \leq \frac{B_1\sqrt{B_1}}{\sqrt{4B_1 + |(3 - \lambda)B_1^2 - 4B_2|}}. \quad (2.16)$$

By subtracting (2.12) from (2.10) and in view of (2.13), we get

$$6(1 + \lambda)a_3 = 6(1 + \lambda)a_2^2 + B_1(b_2 - c_2). \quad (2.17)$$

From (2.4), (2.9), (2.13) and (2.17), it follows that

$$\begin{aligned} |a_3| &\leq |a_2|^2 + \frac{B_1}{6(1 + \lambda)}(|b_2| + |c_2|) \\ &\leq |a_2|^2 + \frac{B_1}{3(1 + \lambda)}(1 - |b_1|^2) \\ &= \left(1 - \frac{4}{3(1 + \lambda)B_1} \right) |a_2|^2 + \frac{B_1}{3(1 + \lambda)}. \end{aligned} \quad (2.18)$$

Substituting (2.16) in (2.18) we obtain the desired inequality (2.2). \square

Remark 2.1. For $\lambda = 0$, the results obtained in the Theorem 2.1 are coincide with results in [24, Theorem 2.1, p.230].

Corollary 2.1. *Let $f \in \mathcal{K}_\sigma(\varphi)$. Then*

$$|a_2| \leq \frac{B_1\sqrt{B_1}}{\sqrt{4B_1 + |2B_1^2 - 4B_2|}} \quad (2.19)$$

and

$$|a_3| \leq \begin{cases} \left(1 - \frac{2}{3B_1}\right) \frac{B_1^3}{4B_1 + |2B_1^2 - 4B_2|} + \frac{B_1}{6} & ; B_1 \geq \frac{2}{3}; \\ \frac{B_1}{3(1+\lambda)} & ; B_1 < \frac{2}{3}. \end{cases} \quad (2.20)$$

3. FEKETE-SZEGÖ INEQUALITIES

In order to derive our result, we shall need the following lemma.

Lemma 3.1. (see [12] or [16]) Let $p(z) = 1 + p_1z + p_2z^2 + \dots \in \mathcal{P}$, where \mathcal{P} is the family of all functions p , analytic in \mathbb{U} , for which $\Re\{p(z)\} > 0$, $z \in \mathbb{U}$. Then

$$|p_n| \leq 2; \quad n = 1, 2, 3, \dots,$$

and

$$\left|p_2 - \frac{1}{2}p_1^2\right| \leq 2 - \frac{1}{2}|p_1|^2.$$

Theorem 3.1. Let f of the form (1.1) be in $\mathcal{G}_\sigma^\lambda(\varphi)$. Then

$$|a_2| \leq \begin{cases} \sqrt{\frac{B_1}{3-\lambda}}, & \text{if } |B_2| \leq B_1; \\ \sqrt{\frac{|B_2|}{3-\lambda}}, & \text{if } |B_2| \geq B_1 \end{cases} \quad (3.1)$$

and

$$\left|a_3 - \frac{4\lambda}{3+3\lambda}a_2^2\right| \leq \begin{cases} \frac{B_1}{3+3\lambda}, & \text{if } |B_2| \leq B_1; \\ \frac{|B_2|}{3+3\lambda}, & \text{if } |B_2| \geq B_1. \end{cases} \quad (3.2)$$

Proof. Since $f \in \mathcal{G}_\sigma^\lambda(\varphi)$, there exist two analytic functions $r, s : \mathbb{U} \rightarrow \mathbb{U}$, with $r(0) = 0 = s(0)$, such that

$$(1 - \lambda)f'(z) + \lambda \left(1 + \frac{zf''(z)}{f'(z)}\right) = \varphi(r(z)) \quad (3.3)$$

and

$$(1 - \lambda)g'(w) + \lambda \left(1 + \frac{wg''(w)}{g'(w)}\right) = \varphi(s(w)). \quad (3.4)$$

Define the functions p and q by

$$p(z) = \frac{1 + r(z)}{1 - r(z)} = 1 + p_1z + p_2z^2 + p_3z^3 + \dots$$

and

$$q(w) = \frac{1 + s(w)}{1 - s(w)} = 1 + q_1w + q_2w^2 + q_3w^3 + \dots$$

or equivalently,

$$r(z) = \frac{p(z) - 1}{p(z) + 1} = \frac{1}{2} \left(p_1z + \left(p_2 - \frac{p_1^2}{2} \right) z^2 + \left(p_3 + \frac{p_1}{2} \left(\frac{p_1^2}{2} - p_2 \right) - \frac{p_1p_2}{2} \right) z^3 + \dots \right) \quad (3.5)$$

and

$$s(w) = \frac{q(w) - 1}{q(w) + 1} = \frac{1}{2} \left(q_1 w + \left(q_2 - \frac{q_1^2}{2} \right) w^2 + \left(q_3 + \frac{q_1}{2} \left(\frac{q_1^2}{2} - q_2 \right) - \frac{q_1 q_2}{2} \right) w^3 + \dots \right). \quad (3.6)$$

Using (3.5) and (3.6) in (3.3) and (3.4), we have

$$(1 - \lambda)f'(z) + \lambda \left(1 + \frac{zf''(z)}{f'(z)} \right) = \varphi \left(\frac{p(z) - 1}{p(z) + 1} \right) \quad (3.7)$$

and

$$(1 - \lambda)g'(w) + \lambda \left(1 + \frac{wg''(w)}{g'(w)} \right) = \varphi \left(\frac{q(w) - 1}{q(w) + 1} \right). \quad (3.8)$$

Again using (3.5) and (3.6) along with (1.2), it is evident that

$$\varphi \left(\frac{p(z) - 1}{p(z) + 1} \right) = 1 + \frac{1}{2} B_1 p_1 z + \left(\frac{1}{2} B_1 \left(p_2 - \frac{1}{2} p_1^2 \right) + \frac{1}{4} B_2 p_1^2 \right) z^2 + \dots \quad (3.9)$$

and

$$\varphi \left(\frac{q(w) - 1}{q(w) + 1} \right) = 1 + \frac{1}{2} B_1 q_1 w + \left(\frac{1}{2} B_1 \left(q_2 - \frac{1}{2} q_1^2 \right) + \frac{1}{4} B_2 q_1^2 \right) w^2 + \dots \quad (3.10)$$

It follows from (3.7), (3.8), (3.9) and (3.10) that

$$\begin{aligned} 2a_2 &= \frac{1}{2} B_1 p_1 \\ 3(1 + \lambda)a_3 - 4\lambda a_2^2 &= \frac{1}{2} B_1 \left(p_2 - \frac{1}{2} p_1^2 \right) + \frac{1}{4} B_2 p_1^2 \end{aligned} \quad (3.11)$$

$$\begin{aligned} -2a_2 &= \frac{1}{2} B_1 q_1 \\ 2(\lambda + 3)a_2^2 - 3(1 + \lambda)a_3 &= \frac{1}{2} B_1 \left(q_2 - \frac{1}{2} q_1^2 \right) + \frac{1}{4} B_2 q_1^2. \end{aligned} \quad (3.12)$$

Dividing (3.11) by $3 + 3\lambda$ and taking the absolute values we obtain

$$\left| a_3 - \frac{4\lambda}{3 + 3\lambda} a_2^2 \right| \leq \frac{B_1}{6 + 6\lambda} \left| p_2 - \frac{1}{2} p_1^2 \right| + \frac{|B_2|}{12 + 12\lambda} |p_1|^2.$$

Now applying Lemma 3.1, we have

$$\left| a_3 - \frac{4\lambda}{3 + 3\lambda} a_2^2 \right| \leq \frac{B_1}{3 + 3\lambda} + \frac{|B_2| - B_1}{12 + 12\lambda} |p_1|^2.$$

Therefore

$$\left| a_3 - \frac{4\lambda}{3 + 3\lambda} a_2^2 \right| \leq \begin{cases} \frac{B_1}{3 + 3\lambda}, & \text{if } |B_2| \leq B_1; \\ \frac{|B_2|}{3 + 3\lambda}, & \text{if } |B_2| \geq B_1. \end{cases}$$

Adding (3.11) and (3.12), we have

$$(6 - 2\lambda)a_2^2 = \frac{B_1}{2}(p_2 + q_2) - \frac{(B_1 - B_2)}{4}(p_1^2 + q_1^2). \quad (3.13)$$

Dividing (3.13) by $6 - 2\lambda$ and taking the absolute values we obtain

$$|a_2|^2 \leq \frac{1}{6 - 2\lambda} \left[\frac{B_1}{2} \left| p_2 - \frac{1}{2} p_1^2 \right| + \frac{|B_2|}{4} |p_1|^2 + \frac{B_1}{2} \left| q_2 - \frac{1}{2} q_1^2 \right| + \frac{|B_2|}{4} |q_1|^2 \right].$$

Once again, apply Lemma 3.1 to obtain

$$|a_2|^2 \leq \frac{1}{6 - 2\lambda} \left[\frac{B_1}{2} \left(2 - \frac{1}{2} |p_1|^2 \right) + \frac{|B_2|}{4} |p_1|^2 + \frac{B_1}{2} \left(2 - \frac{1}{2} |q_1|^2 \right) + \frac{|B_2|}{4} |q_1|^2 \right].$$

Upon simplification we obtain

$$|a_2|^2 \leq \frac{1}{6 - 2\lambda} \left[2B_1 + \frac{|B_2| - B_1}{2} (|p_1|^2 + |q_1|^2) \right].$$

Therefore

$$|a_2| \leq \begin{cases} \sqrt{\frac{B_1}{3-\lambda}}, & \text{if } |B_2| \leq B_1; \\ \sqrt{\frac{|B_2|}{3-\lambda}}, & \text{if } |B_2| \geq B_1 \end{cases}$$

which completes the proof. \square

Remark 3.1. Taking

$$\varphi(z) = \left(\frac{1+z}{1-z} \right)^\beta = 1 + 2\beta z + 2\beta^2 z^2 + \dots, \quad 0 < \beta \leq 1 \quad (3.14)$$

the inequalities (3.1) and (3.2) become

$$|a_2| \leq \sqrt{\frac{2\beta}{3-\lambda}} \quad \text{and} \quad \left| a_3 - \frac{4\lambda}{3+3\lambda} a_2^2 \right| \leq \frac{2\beta}{3+3\lambda}. \quad (3.15)$$

For

$$\varphi(z) = \frac{1 + (1-2\beta)z}{1-z} = 1 + 2(1-\beta)z + 2(1-\beta)z^2 + \dots, \quad 0 \leq \beta < 1 \quad (3.16)$$

the inequalities (3.1) and (3.2) become

$$|a_2| \leq \sqrt{\frac{2(1-\beta)}{3-\lambda}} \quad \text{and} \quad \left| a_3 - \frac{4\lambda}{3-\lambda} a_2^2 \right| \leq \frac{2(1-\beta)}{3+3\lambda}. \quad (3.17)$$

4. BOUNDS FOR THE SECOND HANKEL DETERMINANT OF $\mathcal{G}_\sigma^\lambda(\beta)$

Next we state the following lemmas to establish the desired bounds in our study.

Lemma 4.1. [25] *If the function $p \in \mathcal{P}$ is given by the series*

$$p(z) = 1 + p_1 z + p_2 z^2 + p_3 z^3 + \dots, \quad (4.1)$$

then the following sharp estimate holds:

$$|p_n| \leq 2, \quad n = 1, 2, \dots. \quad (4.2)$$

Lemma 4.2. [15] *If the function $p \in \mathcal{P}$ is given by the series (4.1), then*

$$\begin{aligned} 2c_2 &= c_1^2 + x(4 - c_1^2) \\ 4c_3 &= c_1^3 + 2c_1(4 - c_1^2)x - c_1(4 - c_1^2)x^2 + 2(4 - c_1^2)(1 - |x|^2)z \end{aligned}$$

for some x, z with $|x| \leq 1$ and $|z| \leq 1$.

The following theorem provides a bound for the second Hankel determinant of the functions in the class $\mathcal{G}_\sigma^\lambda(\beta)$.

Theorem 4.1. *Let f of the form (1.1) be in $\mathcal{G}_\sigma^\lambda(\beta)$. Then*

$$|a_2a_4 - a_3^2| \leq \begin{cases} \frac{(1-\beta)^2}{2(1+2\lambda)} [(2-\lambda)(1-\beta)^2 + 1] ; \\ \beta \in \left[0, 1 - \frac{(1+2\lambda) + \sqrt{(1+2\lambda)^2 + 18(1+\lambda)^2(2-\lambda)}}{6(1+\lambda)(2-\lambda)} \right] \\ \frac{(1-\beta)^2}{72(1+2\lambda)} \left(\frac{36[8(1+2\lambda)(2-\lambda) - (1+2\lambda)^2](1-\beta)^2}{-324(1+\lambda)(1+2\lambda)(1-\beta) + 288(1+2\lambda) - 729(1+\lambda)^2} \right. \\ \left. \frac{+8(1+2\lambda) - 18(1+\lambda)^2}{9(1+\lambda)^2(2-\lambda)(1-\beta)^2 - 6(1+\lambda)(1+2\lambda)(1-\beta)} \right) ; \\ \beta \in \left(1 - \frac{(1+2\lambda) + \sqrt{(1+2\lambda)^2 + 18(1+\lambda)^2(2-\lambda)}}{6(1+\lambda)(2-\lambda)}, 1 \right) . \end{cases}$$

Proof. Let $f \in \mathcal{G}_\sigma^\lambda(\beta)$. Then

$$(1-\lambda)f'(z) + \lambda \left(1 + \frac{zf''(z)}{f'(z)} \right) = \beta + (1-\beta)p(z) \quad (4.3)$$

and

$$(1-\lambda)g'(w) + \lambda \left(1 + \frac{wg''(w)}{g'(w)} \right) = \beta + (1-\beta)q(w), \quad (4.4)$$

where $p, q \in \mathcal{P}$ and defined by

$$p(z) = 1 + c_1z + c_2z^2 + c_3z^3 + \dots \quad (4.5)$$

and

$$q(z) = 1 + d_1w + d_2w^2 + d_3w^3 + \dots \quad (4.6)$$

It follows from (4.3), (4.4), (4.5) and (4.6) that

$$2a_2 = (1-\beta)c_1 \quad (4.7)$$

$$3(1+\lambda)a_3 - 4\lambda a_2^2 = (1-\beta)c_2 \quad (4.8)$$

$$4(1+2\lambda)a_4 - 18\lambda a_2a_3 + 8\lambda a_2^3 = (1-\beta)c_3 \quad (4.9)$$

and

$$-2a_2 = (1-\beta)d_1 \quad (4.10)$$

$$2(3+\lambda)a_2^2 - 3(1+\lambda)a_3 = (1-\beta)d_2 \quad (4.11)$$

$$2(10+11\lambda)a_2a_3 - 4(5+3\lambda)a_2^3 - 4(1+2\lambda)a_4 = (1-\beta)d_3. \quad (4.12)$$

From (4.7) and (4.10), we find that

$$c_1 = -d_1 \quad (4.13)$$

and

$$a_2 = \frac{1-\beta}{2}c_1. \quad (4.14)$$

Now, from (4.8), (4.11) and (4.14), we have

$$a_3 = \frac{(1-\beta)^2}{4}c_1^2 + \frac{1-\beta}{6(1+\lambda)}(c_2 - d_2). \quad (4.15)$$

Also, from (4.9) and (4.12), we find that

$$a_4 = \frac{5\lambda(1-\beta)^3}{16(1+2\lambda)}c_1^3 + \frac{5(1-\beta)^2}{24(1+\lambda)}c_1(c_2 - d_2) + \frac{1-\beta}{8(1+2\lambda)}(c_3 - d_3). \quad (4.16)$$

Then, we can establish that

$$\begin{aligned} |a_2a_4 - a_3^2| &= \left| \frac{(\lambda-2)(1-\beta)^4}{32(1+2\lambda)}c_1^4 + \frac{(1-\beta)^3}{48(1+\lambda)}c_1^2(c_2 - d_2) \right. \\ &\quad \left. + \frac{(1-\beta)^2}{16(1+2\lambda)}c_1(c_3 - d_3) - \frac{(1-\beta)^2}{36(1+\lambda)^2}(c_2 - d_2)^2 \right|. \end{aligned} \quad (4.17)$$

According to Lemma 4.2 and (4.13), we write

$$c_2 - d_2 = \frac{(4 - c_1^2)}{2}(x - y) \quad (4.18)$$

$$\begin{aligned} c_3 - d_3 &= \frac{c_1^3}{2} + \frac{c_1(4 - c_1^2)(x + y)}{2} - \frac{c_1(4 - c_1^2)(x^2 + y^2)}{4} \\ &\quad + \frac{(4 - c_1^2)[(1 - |x|^2)z - (1 - |y|^2)w]}{2} \end{aligned} \quad (4.19)$$

for some x, y, z and w with $|x| \leq 1$, $|y| \leq 1$, $|z| \leq 1$ and $|w| \leq 1$. Using (4.18) and (4.19) in (4.17), we have

$$\begin{aligned} |a_2a_4 - a_3^2| &= \left| \frac{(\lambda-2)(1-\beta)^4}{32(1+2\lambda)}c_1^4 + \frac{(1-\beta)^3}{96(1+\lambda)}c_1^2(4 - c_1^2)(x - y) + \frac{(1-\beta)^2}{16(1+2\lambda)}c_1 \right. \\ &\quad \times \left[\frac{c_1^3}{2} + \frac{c_1(4 - c_1^2)(x + y)}{2} - \frac{c_1(4 - c_1^2)(x^2 + y^2)}{4} \right. \\ &\quad \left. \left. + \frac{(4 - c_1^2)[(1 - |x|^2)z - (1 - |y|^2)w]}{2} \right] - \frac{(1-\beta)^2(4 - c_1^2)^2}{144(1+\lambda)^2}(x - y)^2 \right| \\ &\leq \frac{(2-\lambda)(1-\beta)^4}{32(1+2\lambda)}c_1^4 + \frac{(1-\beta)^2c_1^4}{32(1+2\lambda)} + \frac{(1-\beta)^2c_1(4 - c_1^2)}{16(1+2\lambda)} \\ &\quad + \left[\frac{(1-\beta)^3}{96(1+\lambda)}c_1^2(4 - c_1^2) + \frac{(1-\beta)^2}{32(1+2\lambda)}c_1^2(4 - c_1^2) \right] (|x| + |y|) \\ &\quad + \left[\frac{(1-\beta)^2}{64(1+2\lambda)}c_1^2(4 - c_1^2) - \frac{(1-\beta)^2}{32(1+2\lambda)}c_1(4 - c_1^2) \right] (|x|^2 + |y|^2) \\ &\quad + \frac{(1-\beta)^2(4 - c_1^2)^2}{144(1+\lambda)^2}(|x| + |y|)^2. \end{aligned}$$

Since $p \in \mathcal{P}$, so $|c_1| \leq 2$. Letting $c_1 = c$, we may assume without restriction that $c \in [0, 2]$. Thus, for $\gamma_1 = |x| \leq 1$ and $\gamma_2 = |y| \leq 1$, we obtain

$$|a_2a_4 - a_3^2| \leq T_1 + T_2(\gamma_1 + \gamma_2) + T_3(\gamma_1^2 + \gamma_2^2) + T_4(\gamma_1 + \gamma_2)^2 = F(\gamma_1, \gamma_2),$$

$$T_1 = T_1(c) = \frac{(2-\lambda)(1-\beta)^4}{32(1+2\lambda)}c^4 + \frac{(1-\beta)^2c^4}{32(1+2\lambda)} + \frac{(1-\beta)^2c(4-c^2)}{16(1+2\lambda)} \geq 0$$

$$T_2 = T_2(c) = \frac{(1-\beta)^3c^2(4-c^2)}{96(1+\lambda)} + \frac{(1-\beta)^2c^2(4-c^2)}{32(1+2\lambda)} \geq 0$$

$$T_3 = T_3(c) = \frac{(1-\beta)^2c^2(4-c^2)}{64(1+2\lambda)} - \frac{(1-\beta)^2c(4-c^2)}{32(1+2\lambda)} \leq 0$$

$$T_4 = T_4(c) = \frac{(1-\beta)^2(4-c^2)^2}{144(1+\lambda)^2} \geq 0.$$

Now we need to maximize $F(\gamma_1, \gamma_2)$ in the closed square $\mathbb{S} := \{(\gamma_1, \gamma_2) : 0 \leq \gamma_1 \leq 1, 0 \leq \gamma_2 \leq 1\}$ for $c \in [0, 2]$. We must investigate the maximum of $F(\gamma_1, \gamma_2)$ according to $c \in (0, 2)$, $c = 0$ and $c = 2$ taking into account the sign of $F_{\gamma_1\gamma_1}F_{\gamma_2\gamma_2} - (F_{\gamma_1\gamma_2})^2$.

Firstly, let $c \in (0, 2)$. Since $T_3 < 0$ and $T_3 + 2T_4 > 0$ for $c \in (0, 2)$, we conclude that

$$F_{\gamma_1\gamma_1}F_{\gamma_2\gamma_2} - (F_{\gamma_1\gamma_2})^2 < 0.$$

Thus, the function F cannot have a local maximum in the interior of the square \mathbb{S} . Now, we investigate the maximum of F on the boundary of the square \mathbb{S} .

For $\gamma_1 = 0$ and $0 \leq \gamma_2 \leq 1$ (similarly $\gamma_2 = 0$ and $0 \leq \gamma_1 \leq 1$) we obtain

$$F(0, \gamma_2) = G(\gamma_2) = T_1 + T_2\gamma_2 + (T_3 + T_4)\gamma_2^2.$$

(i) The case $T_3 + T_4 \geq 0$: In this case for $0 < \gamma_2 < 1$ and any fixed c with $0 < c < 2$, it is clear that $G'(\gamma_2) = 2(T_3 + T_4)\gamma_2 + T_2 > 0$, that is, $G(\gamma_2)$ is an increasing function. Hence, for fixed $c \in (0, 2)$, the maximum of $G(\gamma_2)$ occurs at $\gamma_2 = 1$ and

$$\max G(\gamma_2) = G(1) = T_1 + T_2 + T_3 + T_4.$$

(ii) The case $T_3 + T_4 < 0$: Since $T_2 + 2(T_3 + T_4) \geq 0$ for $0 < \gamma_2 < 1$ and any fixed c with $0 < c < 2$, it is clear that $T_2 + 2(T_3 + T_4) < 2(T_3 + T_4)\gamma_2 + T_2 < T_2$ and so $G'(\gamma_2) > 0$. Hence for fixed $c \in (0, 2)$, the maximum of $G(\gamma_2)$ occurs at $\gamma_2 = 1$ and also for $c = 2$ we obtain

$$F(\gamma_1, \gamma_2) = \frac{(1-\beta)^2}{2(1+2\lambda)} [(2-\lambda)(1-\beta)^2 + 1]. \quad (4.20)$$

Taking into account the value (4.20) and the cases *i* and *ii*, for $0 \leq \gamma_2 < 1$ and any fixed c with $0 \leq c \leq 2$ we have

$$\max G(\gamma_2) = G(1) = T_1 + T_2 + T_3 + T_4.$$

For $\gamma_1 = 1$ and $0 \leq \gamma_2 \leq 1$ (similarly $\gamma_2 = 1$ and $0 \leq \gamma_1 \leq 1$), we obtain

$$F(1, \gamma_2) = H(\gamma_2) = (T_3 + T_4)\gamma_2^2 + (T_2 + 2T_4)\gamma_2 + T_1 + T_2 + T_3 + T_4.$$

Similarly, to the above cases of $T_3 + T_4$, we get that

$$\max H(\gamma_2) = H(1) = T_1 + 2T_2 + 2T_3 + 4T_4.$$

Since $G(1) \leq H(1)$ for $c \in (0, 2)$, $\max F(\gamma_1, \gamma_2) = F(1, 1)$ on the boundary of the square \mathbb{S} . Thus the maximum of F occurs at $\gamma_1 = 1$ and $\gamma_2 = 1$ in the closed square \mathbb{S} .

Let $K : (0, 2) \rightarrow \mathbb{R}$

$$K(c) = \max F(\gamma_1, \gamma_2) = F(1, 1) = T_1 + 2T_2 + 2T_3 + 4T_4. \quad (4.21)$$

Substituting the values of T_1, T_2, T_3 and T_4 in the function K defined by (4.21), yields

$$\begin{aligned} K(c) = \frac{(1-\beta)^2}{288(1+\lambda)^2(1+2\lambda)} \{ & [9(1-\beta)^2(1+\lambda)^2(2-\lambda) \\ & - 6(1-\beta)(1+\lambda)(1+2\lambda) - 18(1+\lambda)^2 + 8(1+2\lambda)] c^4 \\ & + [24(1-\beta)(1+\lambda)(1+2\lambda) + 108(1+\lambda)^2 - 64(1+2\lambda)] c^2 \\ & + 128(1+2\lambda) \}. \end{aligned}$$

Assume that $K(c)$ has a maximum value in an interior of $c \in (0, 2)$, by elementary calculation, we find

$$\begin{aligned} K'(c) = \frac{(1-\beta)^2}{72(1+\lambda)^2(1+2\lambda)} \{ & [9(1-\beta)^2(1+\lambda)^2(2-\lambda) \\ & - 6(1-\beta)(1+\lambda)(1+2\lambda) - 18(1+\lambda)^2 + 8(1+2\lambda)] c^3 \\ & + [12(1-\beta)(1+\lambda)(1+2\lambda) + 54(1+\lambda)^2 - 32(1+2\lambda)] c \}. \end{aligned}$$

After some calculations we concluded the following cases:

Case 4.1. Let

$$[9(1-\beta)^2(1+\lambda)^2(2-\lambda) - 6(1-\beta)(1+\lambda)(1+2\lambda) - 18(1+\lambda)^2 + 8(1+2\lambda)] \geq 0,$$

that is,

$$\beta \in \left[0, 1 - \frac{(1+2\lambda) + \sqrt{(1+2\lambda)^2 + (2-\lambda)[18(1+\lambda)^2 - 8(1+2\lambda)]}}{3(1+\lambda)(2-\lambda)} \right].$$

Therefore $K'(c) > 0$ for $c \in (0, 2)$. Since K is an increasing function in the interval $(0, 2)$, maximum point of K must be on the boundary of $c \in [0, 2]$, that is, $c = 2$. Thus, we have

$$\max_{0 < c < 2} K(c) = K(2) = \frac{(1-\beta)^2}{2(1+2\lambda)} [(2-\lambda)(1-\beta)^2 + 1].$$

Case 4.2. Let

$$[9(1-\beta)^2(1+\lambda)^2(2-\lambda) - 6(1-\beta)(1+\lambda)(1+2\lambda) - 18(1+\lambda)^2 + 8(1+2\lambda)] < 0,$$

that is,

$$\beta \in \left[1 - \frac{(1+2\lambda) + \sqrt{(1+2\lambda)^2 + (2-\lambda)[18(1+\lambda)^2 - 8(1+2\lambda)]}}{3(1+\lambda)(2-\lambda)}, 1 \right].$$

Then $K'(c) = 0$ implies the real critical point $c_{01} = 0$ or

$$c_{02} = \sqrt{\frac{-12(1+\lambda)(1+2\lambda)(1-\beta) - 54(1+\lambda)^2 + 32(1+2\lambda)}{9(1-\beta)^2(1+\lambda)^2(2-\lambda) - 6(1-\beta)(1+\lambda)(1+2\lambda) - 18(1+\lambda)^2 + 8(1+2\lambda)}}.$$

When

$$\beta \in \left(1 - \frac{(1+2\lambda) + \sqrt{(1+2\lambda)^2 + (2-\lambda)[18(1+\lambda)^2 - 8(1+2\lambda)]}}{3(1+\lambda)(2-\lambda)}, 1 - \frac{(1+2\lambda) + \sqrt{(1+2\lambda)^2 + 18(1+\lambda)^2(2-\lambda)}}{6(1+\lambda)(2-\lambda)} \right].$$

We observe that $c_{0_2} \geq 2$, that is, c_{0_2} is out of the interval $(0, 2)$. Therefore, the maximum value of $K(c)$ occurs at $c_{0_1} = 0$ or $c = c_{0_2}$ which contradicts our assumption of having the maximum value at the interior point of $c \in [0, 2]$. Since K is an increasing function in the interval $(0, 2)$, maximum point of K must be on the boundary of $c \in [0, 2]$ that is $c = 2$. Thus, we have

$$\max_{0 \leq c \leq 2} K(c) = K(2) = \frac{(1-\beta)^2}{2(1+2\lambda)} [1 + (2-\lambda)(1-\beta)^2].$$

When $\beta \in \left(1 - \frac{(1+2\lambda) + \sqrt{(1+2\lambda)^2 + 18(1+\lambda)^2(2-\lambda)}}{6(1+\lambda)(2-\lambda)}, 1\right)$, we observe that $c_{0_2} < 2$, that is, c_{0_2} is an interior of the interval $[0, 2]$. Since $K''(c_{0_2}) < 0$, the maximum value of $K(c)$ occurs at $c = c_{0_2}$. Thus, we have

$$\begin{aligned} \max_{0 \leq c \leq 2} K(c) &= K(c_{0_2}) \\ &= \frac{(1-\beta)^2}{72(1+2\lambda)} \left(\frac{36[8(1+2\lambda)(2-\lambda) - (1+2\lambda)^2](1-\beta)^2}{9(1+\lambda)^2(2-\lambda)(1-\beta)^2} \right. \\ &\quad \left. - \frac{-324(1+\lambda)(1+2\lambda)(1-\beta) + 288(1+2\lambda) - 729(1+\lambda)^2}{-6(1+\lambda)(1+2\lambda)(1-\beta) + 8(1+2\lambda) - 18(1+\lambda)^2} \right). \end{aligned}$$

This completes the proof. \square

Corollary 4.1. *Let f of the form (1.1) be in \mathcal{H}_σ^β . Then*

$$|a_2a_4 - a_3^2| \leq \begin{cases} \frac{(1-\beta)^2[1+2(1-\beta)^2]}{2} ; & \beta \in \left[0, \frac{11-\sqrt{37}}{12}\right] \\ \frac{(1-\beta)^2[60\beta^2-84\beta-25]}{16(9\beta^2-15\beta+1)} ; & \beta \in \left(\frac{11-\sqrt{37}}{12}, 1\right). \end{cases}$$

Corollary 4.2. *Let f of the form (1.1) be in $\mathcal{K}_\sigma(\beta)$. Then*

$$|a_2a_4 - a_3^2| \leq \frac{(1-\beta)^2}{24} \left[\frac{5\beta^2 + 8\beta - 32}{3\beta^2 - 3\beta - 4} \right].$$

Corollary 4.3. *Let f of the form (1.1) be in \mathcal{H}_σ . Then*

$$|a_2a_4 - a_3^2| \leq \frac{3}{2}.$$

Corollary 4.4. *Let f of the form (1.1) be in \mathcal{K}_σ . Then*

$$|a_2a_4 - a_3^2| \leq \frac{1}{3}.$$

REFERENCES

- [1] H. S. Al-Amiri and M. O. Reade, On a linear combination of some expressions in the theory of the univalent functions, *Monatsh. Math.*, **80** (1975), no. 4, 257–264.
- [2] R. M. Ali, S. K. Lee, V. Ravichandran and S. Supramaniam, The Fekete-Szegő coefficient functional for transforms of analytic functions, *Bull. Iranian Math. Soc.*, **35** (2009), no. 2, 119–142, 276.
- [3] R. M. Ali, S. K. Lee, V. Ravichandran and S. Supramaniam, Coefficient estimates for bi-univalent Ma-Minda starlike and convex functions, *Appl. Math. Lett.*, **25** (2012), no. 3, 344–351.
- [4] S. Azizi, A. Ebadian, Sh. Najafzadeh, Coefficient estimates for a subclass of bi-univalent functions, *Commun. Adv. Comput. Sci. Appl.*, **1** (2015), 41–44.

- [5] D. A. Brannan, T. S. Taha, On some classes of bi-univalent functions, *Studia Univ. Babes-Bolyai Math.*, **31**(1986), no. 2, 70–77.
- [6] S. Bulut, Coefficient estimates for a class of analytic and bi-univalent functions, *Novi Sad J. Math.*, **43** (2013), no. 2, 59–65.
- [7] M. Çağlar, H. Orhan and N. Yağmur, Coefficient bounds for new subclasses of bi-univalent functions, *Filomat*, **27** (2013), no. 7, 1165–1171.
- [8] D. G. Cantor, Power series with integral coefficients, *Bull. Amer. Math. Soc.*, **69** (1963), 362–366.
- [9] V. K. Deekonda and R. Thoutreedy, An upper bound to the second Hankel determinant for functions in Mocanu class, *Vietnam J. Math.*, April-2014 (On line version).
- [10] E. Deniz, M. Çağlar and H. Orhan, Second Hankel determinant for bi-starlike and bi-convex functions of order β , arXiv:1501.01682v1.
- [11] P. Dienes, *The Taylor series: an introduction to the theory of functions of a complex variable*, Dover, New York, 1957.
- [12] P. L. Duren, *Univalent functions*, Grundlehren der Mathematischen Wissenschaften, 259, Springer, New York, 1983.
- [13] M. Fekete and G. Szegő, Eine Bemerkung Über Ungerade Schlichte Funktionen, *J. London Math. Soc.*, **S1-8** (1933), no. 2, 85–89.
- [14] B. A. Frasin and M. K. Aouf, New subclasses of bi-univalent functions, *Appl. Math. Lett.*, **24** (2011), no. 9, 1569–1573.
- [15] U. Grenander and G. Szegő, *Toeplitz forms and their applications*, California Monographs in Mathematical Sciences, Univ. California Press, Berkeley, 1958.
- [16] M. Jahangiri, On the coefficients of powers of a class of Bazilevic functions, *Indian J. Pure Appl. Math.*, **17** (1986), no. 9, 1140–1144.
- [17] J. M. Jahangiri, N. Magesh and J. Yamini, Fekete-Szegő inequalities for classes of bi-starlike and bi-convex functions, *Electron. J. Math. Anal. Appl.*, **3** (2015), no. 1, 133–140.
- [18] F. R. Keogh and E. P. Merkes, A coefficient inequality for certain classes of analytic functions, *Proc. Amer. Math. Soc.*, **20** (1969), 8–12.
- [19] K. Lee, V. Ravichandran and S. Supramaniam, Bounds for the second Hankel determinant of certain univalent functions, *J. Inequal. Appl.*, **281** (2013), 1–17.
- [20] W. C. Ma and D. Minda, A unified treatment of some special classes of univalent functions, in: *Proceedings of the Conference on Complex Analysis*, Tianjin, 1992, 157–169, Conf. Proc. Lecture Notes Anal. I, Int. Press, Cambridge, MA, 1994.
- [21] G. Murugusundaramoorthy and N. Magesh, Coefficient inequalities for certain classes of analytic functions associated with Hankel determinant, *Bull. Math. Anal. Appl.*, **1** (2009), no. 3, 85–89.
- [22] H. Orhan, N. Magesh and V. K. Balaji, Initial coefficient bounds for a general class of bi-univalent functions, *Filomat*, **29** (2015), no. 6, 1259–1267.
- [23] H. Orhan, N. Magesh and V. K. Balaji, Fekete-Szegő problem for certain classes of Ma-Minda bi-univalent functions, arXiv:1404.0895v1.
- [24] Z. Peng, Q. Han, On the coefficients of several classes of bi-univalent functions, *Acta Mathematica Scientia*, **34B** (2014), no. 1, 228–240.
- [25] C. Pommerenke, *Univalent functions*, Vandenhoeck & Ruprecht, Göttingen, 1975.
- [26] S. Singh, S. Gupta and S. Singh, On a problem of univalence of functions satisfying a differential inequality, *Math. Inequal. Appl.*, **10** (2007), no. 1, 95–98.
- [27] V. Singh, S. Singh and S. Gupta, A problem in the theory of univalent functions, *Integral Transforms Spec. Funct.*, **16** (2005), no. 2, 179–186.
- [28] H. M. Srivastava, A. K. Mishra and P. Gochhayat, Certain subclasses of analytic and bi-univalent functions, *Appl. Math. Lett.*, **23** (2010), no. 10, 1188–1192.
- [29] P. Zaprawa, On the Fekete-Szegő problem for classes of bi-univalent functions, *Bull. Belg. Math. Soc. Simon Stevin*, **21** (2014), no. 1, 169–178.
- [30] P. Zaprawa, Estimates of initial coefficients for bi-univalent functions, *Abstr. Appl. Anal.*, **2014**, Art. ID 357480, 1–6.

POST-GRADUATE AND RESEARCH DEPARTMENT OF MATHEMATICS
GOVERNMENT ARTS COLLEGE FOR MEN
KRISHNAGIRI 635001, TAMILNADU, INDIA.
E-MAIL ADDRESS: *nmagi_2000@yahoo.co.in*

DEPARTMENT OF MATHEMATICS,
GOVERNMENT FIRST GRADE COLLEGE
VIJAYANAGAR, BANGALORE-560104, KARNATAKA, INDIA.
E-MAIL ADDRESS: *yaminibalaaji@gmail.com*